

UNCLASSIFIED

AD NUMBER
ADB258877
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary information; Jul 1999. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, MD, 21702-5012.
AUTHORITY
USAMRMC ltr, 5 Mar 2002

THIS PAGE IS UNCLASSIFIED

AD _____

GRANT NUMBER DAMD17-98-1-8078

TITLE: Functional Significance of Transcriptional Regulation by
VEGF Receptor Tyrosine Kinases

PRINCIPAL INVESTIGATOR: Adrienne L. Wong

CONTRACTING ORGANIZATION: Duke University Medical Center
Durham, North Carolina 27710

REPORT DATE: July 1999

TYPE OF REPORT: Annual

PREPARED FOR: Commanding General
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government
agencies only (proprietary information, Jul 99). Other requests
for this document shall be referred to U.S. Army Medical Research
and Materiel Command, 504 Scott Street, Fort Detrick, Maryland
21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

DTIC QUALITY INSPECTED 4

20001019 112

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.


LIMITED RIGHTS LEGEND


Award Number: DAMD17-98-1-8078

Organization: Duke University Medical Center

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.





Adrianne L. Wong
Dept of Medicine
435 Sands Building
Duke University Medical Center
Research Drive
Durham, NC 27710-3623

Duke University Medical Center
DURHAM, NORTH CAROLINA
27710

Subject: Unpublished data in Annual Summary Report for Award Number DAMD17-98-1-8078


Commander
U.S. Army Medical Research and Materiel Command
ATTN: MCMR-RMI-S
504 Scott St
Fort Detrick, MD 21702-5012

28 July, 1999

Dear Commander,

The Annual Summary Report for Award Number DAMD17-98-1-8078, "Functional Significance of Transcriptional Regulation by VEGF Receptor Tyrosine Kinases" contains proprietary, unpublished data. Please amend the distribution statement to indicate the limitations of the report. All pages containing proprietary or unpublished data in this report are marked and should be protected as such. Thank-you.

Sincerely,



Adrianne L. Wong

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 1999		3. REPORT TYPE AND DATES COVERED Annual (1 Jul 98 - 30 Jun 99)
4. TITLE AND SUBTITLE Functional Significance of Transcriptional Regulation by VEGF Receptor Tyrosine Kinases				5. FUNDING NUMBERS DAMD17-98-1-8078
6. AUTHOR(S) Adrianne L. Wong				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Duke University Medical Center Durham, North Carolina 27710				8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSORING / MONITORING AGENCY REPORT NUMBER
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Jul 99). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				12b. DISTRIBUTION CODE
13. ABSTRACT. (Maximum 200 words) The relationship between angiogenesis and breast cancer tumorigenesis has been well established. VEGF is a leading candidate for a molecular mediator of tumor angiogenesis. VEGF is expressed by tumor cells in vitro and in vivo; blocking VEGF signaling arrests tumor angiogenesis and tumor growth in vivo. Based on these findings, it is clear that understanding the molecular mechanisms that drive the endothelial responses to VEGF will improve our understanding of tumor angiogenesis. Several of the signaling molecules that associate with the activated VEGF receptors influence pathways involved in transcriptional regulation in the nucleus. We hypothesize that the angiogenic effects of VEGF signaling are attributable, at least in part, to the activation of transcriptional regulation of gene expression. Thus, the goal of the study undertaken is to identify genes that are regulated by VEGF RTK activation and to determine the functional significance of these genes in tumor angiogenesis. We have completed two screens, a cDNA Representational Difference Analysis screen and another using the Gene Discovery Array filters. We further explored the role of C3G and CD9 in VEGF-mediated angiogenesis, but ruled them out. We are currently exploring the role of SOCS2, a negative regulatory protein, in the VEGF angiogenic pathway.				
14. SUBJECT TERMS Angiogenesis, Vascular Endothelium, Receptor Tyrosine Kinase VEGF, Tie, Transcriptional Activation Angiogenesis				15. NUMBER OF PAGES 11
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Limited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

AN Where copyrighted material is quoted, permission has been obtained to use such material.

AN Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

AN Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


NA In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

✓ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

NA In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

NA In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

NA In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


PI - Signature

7/28/99

Date

Table of Contents

Front Cover	1
Letter Regarding distribution statement: unpublished/proprietary data	2
Standard Form 298, Report Documentation Page	3
Foreword	4
Table of Contents	5
Introduction	6
Body	7-9
Appendix	10

Introduction

The relationship between angiogenesis and breast cancer tumorigenesis has been well established. Tumors require angiogenesis to grow and metastasize; the higher the vascular density of a tumor biopsy, the poorer the prognosis. Currently, VEGF is a leading candidate for a molecular mediator of tumor angiogenesis. VEGF is expressed by tumor cells in vitro and in vivo; blocking VEGF signaling arrests tumor angiogenesis and tumor growth in vivo. Based on these findings, it is clear that understanding the molecular mechanisms that drive the endothelial responses to VEGF will improve our understanding of tumor angiogenesis. Several of the signaling molecules that associate with the activated VEGF receptors influence pathways involved in transcriptional regulation in the nucleus. We hypothesize that the angiogenic effects of VEGF signaling are attributable, at least in part, to the activation of transcriptional regulation of gene expression. Thus, the goal of the study undertaken is to identify genes that are regulated by VEGF RTK activation and to determine the functional significance of these genes in tumor angiogenesis. It is hoped that genes regulated by VEGF RTK activation will be useful as biomarkers of tumor angiogenesis or as targets for novel antiangiogenic therapeutics.

Annual Summary

Specific Aim 1: Establish an in vitro system in which to study VEGF RTKs

We have utilized two different endothelial cell lines in our experiments: ECRF (immortalized HUVEC) and HUVEC (primary Human Umbilical Vein Endothelial Cells). Presence and activation of VEGF and Tie receptors was assayed by Western blotting of immunoprecipitated receptors with antiphosphotyrosine antibodies. As a guideline, the maximal expression of "late" genes induced by FGF occurs between 8 and 18 hours. The two conditions used for screening were ECRF cells at 12 hours post-VEGF stimulation for "late" genes and HUVEC at 6 hours post-stimulation for "early" transcriptionally regulated genes.

Specific Aim 2: Clone genes differentially expressed in the endothelium as a result of VEGF RTK activation

cDNA Representational Difference Analysis (RDA) is a PCR-based subtraction approach to find transcriptionally regulated genes. Briefly, mRNA from VEGF-stimulated and unstimulated populations of ECRF cells was reverse-transcribed into cDNAs, digested into short "amplicons", and put through several rounds of PCR-driven subtraction to find the differences between the two populations. RDA yielded 3 putative upregulated gene products and 5 putative downregulated gene products. Contrary to previously published RDA results, these major bands were in fact a heterogeneous population. These putative regulated gene products were subcloned, and a total of 70 genes were sequenced, BLAST searched, and identified. Northern blotting of RNA isolated under the original conditions utilizing the cloned candidate genes as probes narrowed the field to two interesting candidates: C3G and CD9/MRP-1.

Another differential cloning approach recommended by my committee was to screen the Gene Discovery Array filters (GDA, Genome Systems). GDA filters contain an array of roughly 18,000 human genes from the IMAGE consortium of cDNA ESTs. ³²P Probes were made from mRNA from 6 hr VEGF-stimulated or unstimulated HUVEC. Each probe was hybridized to a replicate filter; hybridization was detected on phosphorimager plates and analyzed with the Genome Discovery Software to assess the differences in gene expression between the two conditions. Of the 18,000 ESTs assayed, only 6 ESTs demonstrated an excess of 5 fold expression difference. Northern Blotting was performed using as probes the ESTs that had been identified as potential differential expressors by the GDA screen. Of the 6 potentially regulated ESTs screened, 3 failed to hybridize to Northern blots with a detectable signal, while the other 3 displayed expression levels that were within 1 fold of baseline expression. Possible reasons for the lack of success with this approach include: 1) the timepoint post-VEGF expression was too early, 2) genes relevant to VEGF signaling are activated in a paracrine fashion undetectable by our single cell type system, or 3) the GDA filters simply did not contain targets for VEGF signaling. Therefore, this approach was uninformative.

Specific Aim 3: Characterize the function of genes differentially expressed in the endothelium after stimulation of VEGF RTKs

C3G is a 145 kDa RAS family guanine nucleotide exchange factor for Rap1 and is activated by the expression and binding of Crk adaptor proteins. Crk adaptor proteins are involved in a variety of signaling cascades such as growth factor receptors, integrins, and

cytokines. Interestingly the knockout of B-raf, a member of this signaling pathway, resulted in a vascular phenotype evocative of the Tie2 knockout. While we pulled C3G out of a 12 hour screen, we decided to also investigate its role immediately post VEGF stimulation, when the bulk of cytoplasmic signaling occurs. Western blotting of ECRF cell lysates stimulated with VEGF for 1, 5, 15 minutes or 12 hours demonstrated similar levels of C3G expression compared to unstimulated controls. However, its role might be to differentially associate with other proteins in response to VEGF signaling, so we pursued that line of questioning. In our experiments, C3G did not Co-IP with its upstream partners CrkL or CrkII either with or without VEGF stimulation, despite the abundance of Crk protein in ECRF lysate. In *in vitro* experiments, there was no association between recombinant GST-VEGF receptor kinase domains and CrkL, CrkII, C3G or Rap1 from ECRF cell lysates. Thus, while C3G is present in the endothelial cell milieu it does not have a role in the VEGF signaling pathway under the conditions addressed.

CD9/MRP-1 is a 24 kDa member of the tetraspanin family of cell surface proteins. It has been demonstrated to play a role in cell migration, matrix adhesion, metastasis, and cellular signaling via association with other tetraspanins and integrins. CD9 is expressed in endothelial cells, vascular smooth muscle cells, platelets, melanoma, and carcinoma. Interestingly, in solid tumors, higher CD9 expression correlates with greater metastatic potential. Thus, CD9 could potentially play a role in VEGF-triggered endothelial cell migration or downstream cellular signaling.

We began with experiments to assess CD9 expression in response to VEGF stimulation. A Northern timecourse performed on total RNA from ECRF cells [fig] revealed a gradual increase of CD9 expression to twofold of baseline by 16 hours post VEGF stimulation. FACS analysis of CD9 cell surface protein expression post-VEGF stimulation revealed a rapid 2 fold increase at 0.5 hr post VEGF treatment. Thus, we hypothesized that VEGF stimulation might trigger the localization of CD9 to the cell surface from a sequestered cytoplasmic location in an event parallel to its gradual transcriptional upregulation.

We assessed the functional role of CD9 protein in VEGF-mediated cell migration via a modified Boyden chamber assay. HUVEC cells were plated on the upper surface of a Transwell chamber (Corning) and we tested their ability to migrate through the membrane towards VEGF in the lower chamber. Here, cell surface CD9 function was disrupted by the addition of Syb-1, a monoclonal anti-CD9 blocking antibody (Dr. C. Boucheix INSERM, France). The addition of Syb-1 resulted in a 50% decrease in VEGF-mediated chemotaxis, as compared to untreated and nonspecific IgG-treated controls. This implicates CD9 in VEGF-mediated cell migration. Thus the working hypothesis was that pretreatment with Syb-1 results in an impediment of VEGF-induced integrin-dependent motility via disruption of tetraspanin/integrin interactions.

To investigate the mechanisms of CD9 cell surface localization from vesicular sequestration, we employed confocal microscopy and timelapse microscopy. Von Willebrand's factor (VWF) was used as a marker for Weibel-Palade (WP) bodies, storage vesicles found in endothelial cells. VEGF stimulation did not result in the fusion of WP bodies with the cell surface as compared to controls. Changes in microfilament organization due to VEGF were tracked with a phalloidin stain as a positive control. Double-staining with CD9 and VWF did not result in significant overlap. Therefore, this suggested that WP bodies do not function as CD9 sequestration vesicles.

Immunocytochemistry experiments viewed by confocal microscopy revealed no obvious differences in total vs cell surface CD9 expression 30 minutes post VEGF stimulation [fig]. This might be explained by a constraint of the system: a two-fold difference in cell surface expression

might not be bright enough to be seen in a single-thickness slice. GFP-tagged versions of CD9 were constructed and transiently transfected into endothelial cells to track the expression and localization of CD9 in real time, which generated similar results. Possible reasons for this observation might be the improper processing or targeting of the GFP tagged protein, a perturbation of the normal cellular responses to VEGF stimulation, or it may well be that the amount of CD9 expression at the cell surface is not quantifiably altered by VEGF stimulation.

In conclusion, CD9 was identified by an RDA screen as a gene transcriptionally regulated by VEGF stimulation of ECRF cells. Its expression increases 2 fold over 16 hours post stimulation by Northern blotting and also 2 fold at 0.5 hr post stimulation by FACS. The addition of a CD9 blocking antibody rendered VEGF-mediated HUVEC migration 50% less efficient. While we could reconcile the somewhat contradictory expression data with the explanation that CD9 was playing an immediate role in migration where VEGF stimulation would drive sequestered stores of CD9 protein to the cell surface, thus necessitating refurbishment by transcriptional activation, we could not find any evidence to support this theory. Thus, we believe that the Syb-1 effect on cell migration is an indirect result of binding CD9 on the cell surface and is not specific to VEGF-mediated cell migration.

Future directions

Upon the completion of two screens for genes transcriptionally regulated by VEGF, I had no positive results. Therefore, it was decided in February 1999 to change direction. My mentor, Dr. Kevin Peters, has left Duke University for a position in industry, though he remains my official thesis advisor. I am continuing my graduate studies in the laboratory of Dr. Chris Kontos, a former fellow trained by Dr. Peters, who is investigating the biology and signaling of endothelial RTKs. My academic committee agreed that pursuing another aspect of VEGF signaling would be preferable to performing another round of screening. While the initial purpose of this proposal was to discover novel genes transcriptionally regulated by VEGF signaling, the project that I am currently involved in is focused on earlier events in the signaling cascade, which nonetheless may play an important role in VEGF-mediated gene regulation. As the VEGF signaling pathways has by no means been completely elucidated, novel discoveries involving any aspect of this angiogenic process will be exciting additions to the vernacular.

SOCS2 was pulled out of a yeast-2 hybrid screen of a human fetal heart library using VEGFR-2 as a bait in our lab. SOCS-2 is an SH2 domain protein that belongs to the SOCS-box family of cytoplasmic signaling proteins, which function as suppressors of cytokine signaling. We have discovered a novel kinase-dependent association between the endothelial RTKs (VEGFR1, VEGFR2, Tie2) and SOCS2 in the 2-hybrid system. We are currently making a His-tagged SOCS2 protein to test for association with the GST-fused kinase domains of these same RTK proteins in vitro. Studies are underway to map the binding site between these proteins using the specific domains of SOCS2: N-terminal, SH2, and SOCS-box. Furthermore, our collaborator T. Willson at WEHI, Melbourne, AU has agreed to provide us with the transfection constructs for SOCS-1, SOCS-2, SOCS-2 K/R and SOCS-3 for in vivo cellular association and specificity experiments. We are also generating an adenovirus for efficient introduction and overexpression of SOCS-2 into endothelial cells to look at downstream signaling mechanisms. Hopefully, these experiments will prove successful and yield the understanding of a novel aspect of VEGF signaling.

Appendix

Key research accomplishments

- Completed RDA screen for genes transcriptionally regulated by VEGF stimulation of ECRF cells
- Learned how to perform migration assays, confocal and video microscopy
- Investigated role of c3G and CD9 in VEGF-stimulated angiogenesis
- Completed GDA screen

Reportable outcomes

- Offered postdoctoral positions by Dr. Susan Fisher, UCSF (Endothelial RTKs & early implantation) and Dr. Linda Giudice, Stanford (RTKs and endometriosis).



DEPARTMENT OF THE ARMY

US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

5 Mar 02

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statements

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for grants. Request the limited distribution statements for Accession Documents listed at enclosure be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl



PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

DISTRIBUTION TO BE CHANGED TO UNLIMITED,
APPROVED FOR PUBLIC RELEASE

ACCESSION DOCUMENT NUMBERS

ADB267943	ADB267947
ADB257308	ADB268439
ADB233733	ADB242952
ADB263445	ADB248375
ADB258807	ADB240661
ADB257354	ADB271166
ADB240907	ADB258706
ADB270764	ADB244250
ADB241926	ADB258773
ADB246527	ADB254490
ADB268422	ADB268189
ADB252915	ADB270793
ADB258877	ADB266144
ADB268117	ADB236672
ADB267884	ADB259031
ADB254260	ADB270765
ADB268296	ADB270785
ADB258930	ADB268113
ADB271098	ADB270791